

Motor Imagery Promising Technique for Rehabilitation of Patients with Parkinson's disease: A Systematic Review

Muhammad Kashif^{1*}, Ashfaq Ahmad², Kiran Bashir³, Maryam Farooq⁴,
Saba Maqbool⁴, Iqra Nadeem⁵

ABSTRACT

Recent research has suggested that motor imagery may be helpful for the Rehabilitation of Parkinson's disease (PD) patients. Using MI, a movement can be imagined without muscles being activated. MI induces plastic changes in the motor cortex, improving motor performance. This systematic review was designed to examine the effects of MI on the physical rehabilitation of people with Parkinson's disease. Five databases were used to identify similar studies using selected keywords. This review assessed the "methodological quality of each randomized control trial study" using the eleven-point Physiotherapy Evidence Database scale, widely used to rate physiotherapy "literature". The benefits of using MI to treat patients with PD were identified in this review. Seven studies were identified; 6 studies reported motor function, two studies worked on bradykinesia, two on balance and two reported impairments in Gait in PD using MI. The included studies had small samples, varied methodological approaches, and varied quality from good to fair. According to the current review, MI provides more significant benefits for Parkinson's patients than conventional physical therapy alone when used along with other therapeutic methods for improving motor function and balance.

KEYWORDS: Rehabilitation, Parkinson's disease, Motor imagery, systematic review.

INTRODUCTION

Motor imagery (MI) is a procedure of mentally practicing a motor task or activating specific muscles without explicit feedback¹. This technique is a motor representation related to the preparation and intention of movements through which a person rehearses or repeats a particular action. It is mainly used in sports training as a mental activity exercise and neurological rehabilitation. Functional neuroimaging studies have depicted that during mental activation, frontal, parietal, primary cortex, basal ganglia, supplementary motor area, and cerebellum areas are mainly stimulated, especially during the execution of movement^{2,3}. The premotor cortex and inferior parietal lobule, related to motor cognition, are stimulated during MI. Movement implementation and core motor learning mechanisms are linked in this way. Imagination and

motor execution are, however, highly dependent on task complexity⁴. MI also provides the basis for brain-computer interface development for physically disabled persons, leading to more precise movement execution. MI can be contradictory, depending on how a task or rehearsal is linked with visual, acoustic, somatosensory and balanced perceptions and sensations⁵.

In rehabilitation, the advantage of MI is that it gives freedom of movement capabilities, no chance of physical harm, limited financial costs, increased compliance, and no equipment requirement. In addition, MI also target many motor (for example, range of motion) and non-motor tasks that are cognitive and sensory (for example, self-confidence, pain and motivation) related to performance. MI technique provides various delivery opportunities to physiotherapists virtually, physically and remotely. So, this technique is more relevant to many PD communities, including underserved and remote ones, thus targeting gaps and future directions documented by previous studies⁶. The primary motor cortex (M1) processes MI tasks and motor actions, as do the premotor, supplementary, and parietal motor cortexes⁷. The goal of MI is to imagine that movements or muscles are being activated or moved without actually moving or starting them. In many neurological disorders affecting motor recognition and execution, MI is highly effective at improving motor skills⁸.

Our hands and eyes mostly move in a coordinated manner in the execution of particular movements

¹Riphah College of Rehabilitation and Allied Health Sciences, Riphah International University, Faisalabad Campus, Pakistan

²University Institute of Physical Therapy, Faculty of Allied Health Sciences, University of Lahore, Lahore, Pakistan

³Islamabad College of Physiotherapy, Margalla Institute of Health Sciences, Islamabad, Pakistan

⁴School of Rehabilitation Sciences, The University of Faisalabad, Faisalabad, Pakistan

⁵Sargodha Medical College, University of Sargodha, Sargodha, Pakistan

Correspondence: kashif.shaffi@gmail.com

doi: 10.22442/jlumhs.2023.01014

Received: 03-02-2023

Revised: 02-05-2023

Accepted: 09-05-2023

Published Online: 10-08-2023



during activities of daily living. The study depicted that the presence of visual data while performing purposeful actions adds to endpoint consistency and accuracy. Eye movements while executing the aimed task⁹. It has been shown that MI training sessions increase performance in terms of speed, performance accuracy, muscle strength, movement dynamics, and motor skill performance¹⁰.

Alterations in MI have also been known in many progressive or acute neurological disorders, such as multiple sclerosis, PD, and cerebrovascular lesions. MI is believed to engage similar neural pathways used in motor tasks and linked kinesthetic, visual and sensory input integration¹¹.


In Parkinson's disease, there is a decline in the level of dopamine in the basal ganglia, which causes motor and non-motor symptoms. Its symptoms may be treated with levodopa therapy, mainly as a highly active symptomatic treatment. The majority of long-term complications are managed with medications. However, certain motor features and complications require an active role in rehabilitation. Patients with PD can benefit from several rehabilitation strategies, including the MI technique, to aid them in learning motor skills¹².

One study stated that action observation is a more feasible and safe procedure for Parkinson's patients to improve Gait and balance and reduce falls¹³. Another study reviewed that imagery therapy and home-based action observation are possible in moderate to mild PD and provide an independent, low cost and flexible approach in addition to conventional neuro-rehabilitation¹⁴. In this systematic review, we aimed to identify and appraise studies that examined the role of MI in the physical rehabilitation of patients with Parkinson's disease.

METHODOLOGY

This systematic review was conducted to study the MI as a promising tool in the physical Rehabilitation of PD by using keywords 'Motor Imagery' or 'Mental Practice' or 'Motor Imagery Training' and Parkinson's disease OR 'Parkinsonism' OR 'Parksonian' on online databases, including Web of Science, Cochrane Library, PubMed, CINHALL and PEDro. One thousand six hundred forty articles were recruited from searches on selected databases. After excluding some articles based on duplication, irrelevant population, study design and outcomes, seven full-text articles were selected to add to this review. The study inquiry for this review was: What role may MI play in the physical Rehabilitation of PD? The PICOS strategy (population, intervention, comparison, outcome measures, study design strategy) was employed in this review. Patients with overt visual, cognitive, or hearing impairment were excluded from the study. The review included MI studies alone or with any other combination, randomized clinical trials, and analyses in English. (Figure I)

Figure I:
PRISMA flow diagram Eligibility and data synthesis



Data extraction and quality assessment

Two reviewers carried out the data search for the localization of studies with the established strategy of the current systematic review. Data extraction included age, gender, sample size, intervention, Hoehn-Yahr scale dosage, outcome, and results of the included studies¹⁵ (Table I). A Physiotherapy Evidence Database (PEDro) scale, widely used to measure the quality of studies in physiotherapy-based research, was used to assess the methodological quality of each RCT study. Inter-rater reliability of PEDro scores for physiotherapy interventions is 'fair' to 'excellent' (interclass correlation coefficients of 0.53 -0.91). Inter-rater reliability of PEDro scale items varies from 'fair' to 'almost perfect' (Kappa 0.36-1.00) in physiotherapy trials. Increasing inter-rater reliability occurs when two or three raters agree on the total PEDro score and the individual PEDro scale items. A PEDro score of 0-3 is considered 'poor', a score of 4-5 is considered 'fair', a score of 6-8 is considered 'good', and a score of 9-10 is considered 'excellent'^{16,17}.

RESULTS

Quality appraisal of studies

In total, seven studies, six rated as Good with Score 6 to 8^{14,18-22}. Only one study²³ scored Fair with a score of 4 according to Pedro's rating scale¹⁶. Random allocation, eligibility criteria and baseline prognostic indicators were reported in all seven studies. Out of 7 studies, only two studies^{19,23} lack information regarding concealed allocation. Only one study²³ was open-blinded. All seven studies reported between-group statistical comparisons. (Table II)

Table I: Data Extraction of included studies

Study	Age Range/ Mean age	Gender	Sample size	Experimental Group	Control Group	Hoehn-Yahr scale	Dosage	Outcome Measures	Results
Sarasso E et al. ²¹	Dual-task group: 63.81±9.23 Dual task +AOT-MI group: 67.51±6.12	Dual-task group:8M/4F Dual task +AOT-MI group: 8M/5F	25	4 gait/balance Exercises by Using observation/imagination	4 gait/ balance Exercises without imagination	01-03	6 weeks, 3 times a week, about 1 hour each session	H&Y scale, MDSUPDRS-II, MDS-UPDRS-III, Mini-BEST, TUG,10-MWT, ABC, PDQ-39 & NFOG-Q	The study's findings reported that Using observation/ imagination improved dual-task mobility and balance in patients with PD.
Santiago et al. ²⁰	CG:61.40±9.05 EG:61.30±9.95	Both genders	20	mental practice and physical practice	Physical practice only	02-03	On the day of the initial assessment, a single training session was conducted, with steps 1, 2, 3, 5 and 7 performed by both groups and steps 4 and 6 performed by the experimental group.	Qualisys Motion Capture & TUG Test	The study found that mental training did not significantly improve TUG scale scores compared to physical training.
Lokhandwala M 2019 ²²	60-75Years	Both genders	30	Physical practice and mental imagery were provided to the experimental group (N=15).	The control group received only physical practice. (N=15)	1.5 - 3	Both groups received 20 therapy sessions five days a week for four weeks.	UPDRS, TUG Test, PDQ-39 and Stroop test	Between the two groups, the performance times from supine to standing and standing to supine were significantly different, with a p-value of 0.00, the TUG test with a p-value of 0.003 and the Stroop test with a p-value of 0.004.
Bek J et al. ¹⁴	Range from 47 to 73 Year	M/F: 9/1	10	The experimental group received ACTION-PD combined AO plus MI intervention for individuals with PD. (N= 6)	The control group continued with their usual treatment for PD. (N = 4)	01-03	6-weeks training period	DextQ-24 , KVIQ, PDQ-39	For people with mild to moderate PD, AO + MI training at home, performed via mobile technology, is feasible. In addition, motor performance was better in the experimental group than in the control group.
Abraham A 2018 ²³	Mean age in the Dynamic Neuro-Cognitive Imagery (DNI) group was 66.4 ± 12.5, and in the control group was 65.1±7.5	M/F: 16/4	20	The experimental group received DNI training (n = 10)	In-home learning and exercise program (control; n = 10)	01-03	Experimental and control groups completed at least 16 hours of training within two weeks.	MoCA, CPF, MIQ-RS, KVIQ -20, & VMIQ-2	In this study, PD patients with DNI training had improved imageability, disease severity, and motor and non-motor functions.
Tamir R 2007 ¹⁹	Experimental group was 67.4±9.7 and in the Control group, 67.4±9.1	M/F: 15/8	23	The experimental group was treated with both imagery and real practice	The control group received only physical exercises.	1.5-03	Exercises for both groups were applied during 1-hour sessions held twice a week for 12 weeks.	UPDRS, Stroop and clock drawing tests	Compared to the control group, the treatment group showed significantly higher gains in the mental and motor subgroups of the UPDRS and cognitive tests. There was an improvement in daily living activities for both groups.
Braun S 2011 ¹⁸	The mean age in the Experimental group was 70±8, and in the Control group was 69±8	M: 47	47	physiotherapy, mental practice and usual care	Physiotherapy, relaxation and usual care	01-04	A six-week intervention period for both groups	VAS, TUG, & 10-MWT	The experimental group improved more than the control group, but not significantly.

Table II: Quality assessment of included studies

Study.	Randomly allocation	Concealed Allocation	Baseline comparability	Participant blinding	Therapist blinding	Assessor Blinding	<15% dropouts	Intention to treat analysis	Between-group difference reported	Point estimate and variability reported	Eligibility criteria	Total Score
Lokhandwala M 2019 ²²	Y	N	Y	N	N	Y	Y	Y	Y	Y	Y	7/10
Braun S 2011 ¹⁸	Y	Y	Y	N	N	Y	Y	Y	Y	Y	Y	8/10
Tamir R 2007 ¹⁹	Y	N	Y	N	N	Y	Y	N	Y	Y	Y	6/10
Abraham A 2018 ²³	Y	N	Y	N	N	N	N	N	Y	Y	Y	4/10
Santiago et al. ²⁰	Y	Y	Y	N	N	Y	Y	N	Y	Y	Y	7/10
Sarasso E et al. ²¹	Y	Y	Y	N	N	Y	Y	Y	Y	Y	Y	8/10
Bek J et al. ¹⁴	Y	Y	Y	N	N	Y	Y	Y	Y	Y	Y	8/10

Sample characteristics

A total of 175 patients with PD participated in 7 studies. Out of 175 patients, the male-to-female ratio was 103:22 in 5 studies^{14,18,19,21,23}; 47 male participants were the maximum number¹⁸, and the minimum number was 8²¹. The maximum number of female participants was 8¹⁹. Bek J et al.¹⁴ researched only one female participant. All studies included patients with stage ≤ 4 , according to Hoehn and Yahr classification (H&Y)¹⁵. The total time duration for treatment ranges from 2 weeks to 12 weeks.

Types of Interventions in experimental groups

Types of interventions used in the experimental group were mental imagery of functional activities^{19,20,22}, Tablet computer App¹⁴, Dynamic neuro-cognitive imagery²³, mental practice of walking, standing up from a chair or the floor, mental imagery of gait tasks¹⁸, Dual task+ Action Observation Task - MI²¹.

Types of interventions in the control group

Types of interventions used in the control group were Dual Task²¹, conventional physical therapy²², Physical practice of Gait²⁰, Physical practice¹⁴, in-home learning²³, callisthenic exercises¹⁹, flexibility, muscular strength and coordination exercises, relaxation exercises¹⁹, Physiotherapy with relaxation therapy¹⁸.

OUTCOMES

Bradykinesia

Of 7 studies, two worked on bradykinesia^{19,22}. Lokhandwala M 2019²² reported the effectiveness of the combination in reducing bradykinesia, fear, and anxiety. The study showed improved time and confidence to perform functional activities through integrated physical and mental practice. UPDRS subsets for the experimental group showed less than significant improvement. This improvement is because MI is essential to cognition, motivation and memory. Experimental group. Tamir and fellows¹⁹ also agreed with these findings and showed significant results in reducing bradykinesia and improving motor function.

Outcome Measures for Bradykinesia

Outcomes measures to assess bradykinesia were the performance of a movement sequence, UPDRS¹⁹, TUG (Timed up and go), ST: TS (Supine to Stand), and ST: TS (Stand to Supine) performance time²².

Motor function

Out of 7 studies, six were conducted to determine the effectiveness of MI on motor function¹⁸⁻²³. According to Amit and coworkers²³, Dynamic neuro-cognitive imagery significantly improved motor function measures, specifically in the TUG manual and 360-degree Turn. Sarasso E et al.²¹ reported that combining Action Observation Imagery-MI with the dual task is more beneficial for reorganizing the brain area, and these effects are long-lasting.

Outcome measures for motor function

Unified Parkinson's Disease Rating Scale (UPDRS)^{19,22}, Single and Dual Timed Up and Go (TUG) test^{18,20,21}, Forward Gait Speed, 6-Minute Walk Test (6MWT), 30-Second Chair Stand, 360° Turn Test and Push and Release Test (PRT)²³ were used as outcome measuring tools for motor function.

Balance

Of 7 studies, two assessed balance as an outcome^{19,21}. Tamir R 2007¹⁹ study also in favor of MI along with physical exercises for the improvement of balance.

Outcome measures for balance

Outcome measuring tools used were the Mini Balance Evaluation Systems Test (MiniBESTest)²¹ and the Activities Balance Confidence Scale (ABC)¹⁹.

Gait

Out of 7 studies, two assessed Gait as an outcome. These studies reported improvement in stance and swing time^{20,21}. The intergroup difference was insignificant in one training session; however, the intragroup difference was significant. Braun S 2011¹⁸ described MI does not affect walking performance. However, patients with stages under three on the H&Y scale showed some improvement.

Outcome measures for Gait

New Freezing of Gait Questionnaire (NFoG-Q)²¹ and Qualisys Motion Capture System²⁰ were utilized as outcome measures.

Other outcomes and outcome measures

Dexterity, quality of life, MI ability, visual and kinesthetic ability, vividness, disease severity, psychological measures, and Activity of daily living (ADLs) were evaluated with balance, Gait, motor function, and bradykinesia. No significant improvement was seen in visual and kinesthetic ability and ADLs. Dexterity, cognition, and quality of life were improved. Tamir and fellows significantly improved the mental subset of the Unified Parkinson's Disease Rating Scale (UPDRS).

These outcomes were assessed by Stroop Test²², Dexterity Questionnaire (DextQ-24)¹⁴, Kinesthetic and Visual Imagery Questionnaire (KVIQ)²³, Parkinson's disease Questionnaire (PDQ-39)²³, Montreal Cognitive Assessment (MoCA)²³, Composite Physical Function Scale (CPF)²³, The Movement Imagery Questionnaire-Revised Second Version (MIQ-RS)²³, Vividness of Movement Imagery Questionnaire-Revised Version (VMIQ-2)²³.

DISCUSSION

Using MI for physical rehabilitation among individuals with PD was the aim of this systematic review. The benefits of MI for motor recovery are especially evident when long-term rehabilitation programs are used. In terms of neurorehabilitation, MI appears to be a promising future technique. Several studies have demonstrated improved skills, strength, and function through MI in various neurological conditions²⁴⁻²⁶.

The MI improves motivation, concentration and attention in patients with neurological disorders²⁷. According to Asavari and colleagues, MI adds value to physical therapy and is a practical way to elicit motor-evoked potentials (MEPs) in patients who cannot receive physical therapy because of severe motor impairments. Despite brain lesions, MI has consistently proven effective in motor rehabilitation through increased treatment or neural stimulation²⁸. Compared to other measurement approaches, neurofeedback demonstrates the highest success rates for motor symptoms in the early stages of development²⁹. Parkinsonism is characterized by bradykinesia, also associated with Amyotrophic Lateral Sclerosis, which causes progressive degeneration of upper and lower motor neurons³⁰. Therefore, Abidi M et al.³¹ suggested that postural control can improve gait reorganization by reducing fear of falling, leading to altered gait patterns and locomotion imagery time. Individuals with PD have difficulty working on computers, such as reduced keyboard speed, hitting accidental keys when typing, and difficulty clicking or controlling a mouse. Another study by Woodrow-Hill 2021³⁴ indicated that MI can be an effective technique for smoothing movement in PD patients.

In their study, Gil-Bermejo-Bernardez-Zerpa A 2021³⁵ showed that MI and its combination with relaxation exercises helped improve motor skills, Gait, balance and quality of life in patients with Multiple Sclerosis. A study by Nicholson and colleagues found that MI training could improve balance and mobility in older adults without neurological conditions. The results of this study suggested that MI training could be an adjunct to standard physiotherapy care in older adults²⁵. MI results in functional reorganization of motor control and executive attention areas when combined with dual-task gait/balance training, and has longer-lasting effects on mobility and balance²¹. Bae et al.³² found that combining MI with balance training significantly improved patients' balance and gait abilities more than balance training alone.

Li R-Q 2017³³ found an increase in walking speed of 40% after MI treatment and significant improvements in step, foot rhythm, and other aspects. Woodrow-Hill 2021³⁴ reported that MI provides short-term Gait benefits when measured using walking speed. Another study by Gil-Bermejo-Bernardez-Zerpa A 2021³⁵ reported that MI enables more demanding or complex motor tasks, improving Gait and mobility in patients with PD. Moreover, it is reported that MI enhances the quality of life through the neurofeedback mechanism³⁶.

Integrating MI with physical therapy treatment may reduce bradykinesia in patients with PD. MI could be an essential component of the cognitive strategies provided to these patients³⁷. Avanzino L 2013³⁸ observed the effects of MI on sequential movements in Parkinson's patients. The synchronization-continuation paradigm assessed the motor timing arrears in sequential movements through MI. Results suggested that PD patients showed discerning deficits in motor timing and motor execution, which can affect motor planning. It was also concluded that Timing deficits could affect MI abilities.

MI techniques are used not only for PD rehabilitation but also for other disorders, as recent studies suggest for other neurological disorders such as cerebrovascular accidents, spinal cord injury and non-neurological disorders such as after knee arthroplasty and amputation^{39,40}. Although MI is very effective in various disorders, its limitations still influence its effectiveness. One necessity of MI is a quiet environment and one's concentration skills, so these factors can affect its effectiveness. Other factors that control its efficacy are longer session duration and age-related discrepancies^{39,41}. Limitations reported in studies were small sample size^{6,19,42}, no follow-up⁶, short duration intervention, uncertainty about the dose and sensitivity of assessment tools¹⁹.

MI in the physical rehabilitation of PD has an essential role in activating the brain's neural network, thus enhancing its overall function. In addition to improving motor function, there are various benefits of MI, such

as its non-invasive nature, no safety issues, no need for equipment and trained staff and can be used in home-based intervention³⁹. The only English literature added to our review was one of the limitations of our study. Besides these benefits, further high-quality clinical trials are required to implement MI as a therapeutic tool in the physical Rehabilitation of PD. Furthermore, more extensive studies should examine how MI practice combines with conventional PT for patients with Parkinson's disease, given the clinical and statistical significance of the findings.

CONCLUSION

Based on the results of this systematic review, MI interventions in PD are likely to have beneficial effects on balance, Gait, and motor functions. More studies with large sample sizes and a sound methodology should be conducted to assess the utility of MI. Moreover, technological advances could benefit the administration of training based on MI, and their application to Parkinson's Rehabilitation could lead to new protocols.

Conflict of Interest: There is no conflict of interest between the authors.

Financial Disclosure / Grant Approval: Research work for this study was not funded by any funding agency.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publically.

AUTHOR CONTRIBUTIONS

Kashif M: Data Interpretation, drafting or critically reviewing its intellectual content, Final approval of the manuscript, agreement to be accountable for all aspects of the research work.

Ahmad A: Interpretation; critically reviewing its intellectual content, Final approval of the manuscript, agreement to be accountable for all aspects of the research work, supervision.

Bashir K: The acquisition, literature review, drafting of the work, final approval of the manuscript, AND the agreement to be accountable for all aspects of the research work.

Farooq M: Design of the work, drafting work, Final approval of the manuscript, and integrity of any part of the work are appropriately investigated and resolved.

Iqbal S: Interpretation of data for the work, revising it critically for important intellectual content, Final approval of the manuscript, and integrity of any part of the work are appropriately investigated and resolved.

Nadeem I: Design of the work, manuscript drafting, and all integrity issues related to the work are investigated and resolved appropriately.

REFERENCES

1. Readman MR, Crawford TJ, Linkenauger SA, Bek J, Poliakoff E. Motor imagery vividness and symptom severity in Parkinson's disease. *J Neuropsychology*. 2023; 17(1): 180-92. doi: 10.1111/jnp.12293.
2. Leiguarda R, Cerquetti D, Tenca E, Merello M. Globus pallidus internus firing rate modification after motor-imagination in three Parkinson's disease patients. *J Neural Transm(Vienna)*. 2009; 116(4): 451-5. doi: 10.1007/s00702-009-0203-3. Epub 2009 Mar 10.
3. Kober SE, Grössinger D, Wood G. Effects of motor imagery and visual neurofeedback on activation in the swallowing network: A real-time fMRI study. *Dysphagia*. 2019; 34(6): 879-95. doi: 10.1007/s00455-019-09985-w. Epub 2019 Feb 15.
4. Mizuguchi N, Kanosue K. Changes in brain activity during action observation and motor imagery: Their relationship with motor learning. *Prog Brain Res*. 2017; 234: 189-204. doi: 10.1016/bs.pbr.2017.08.008. Epub 2017 Sep 14.
5. Hanakawa T. Organizing motor imageries. *Neurosci Res*. 2016; 104: 56-63. doi: 10.1016/j.neures.2015.11.003. Epub 2015 Nov 19.
6. Abraham A, Duncan RP, Earhart GM. The Role of Mental Imagery in Parkinson's Disease Rehabilitation. *Brain Sci*. 2021; 11(2): 185. doi: 10.3390/brainsci11020185.
7. Lindsay RS, Larkin P, Kittel A, Spittle M. Mental imagery training programs for developing sport-specific motor skills: a systematic review and meta-analysis. *Phys Educ Sport Pedagogy*. 2023; 28(4): 444-65. doi: 10.1080/17408989.2021.19911297.
8. Castillo-Rangel C, Marin G, Hernández-Contreras KA, Vichi-Ramírez MM, Zarate-Calderon C, Torres-Pineda O et al. Neuroinflammation in Parkinson's Disease: From Gene to Clinic: A Systematic Review. *Int J Mol Sci*. 2023; 24(6): 5792. doi: 10.3390/ijms24065792.
9. Heremans E, Smits-Engelsman B, Caeyenberghs K, Vercruyssen S, Nieuwboer A, Feys P et al. Keeping an eye on imagery: the role of eye movements during motor imagery training. *Neurosci*. 2011; 195: 37-44. doi: 10.1016/j.neuroscience.2011.07.030.
10. Dickstein R, Deutsch JE. Motor imagery in physical therapist practice. *Phys Ther*. 2007; 87(7): 942-53. doi: 10.2522/ptj.20060331. Epub 2007 May 1.
11. Monaco MRL, Laudisio A, Fusco D, Vetrano DL, Ricciardi D, Zuccalà G et al. Laterality in Parkinson's disease may predict motor and visual imagery abilities. *Funct Neurol*. 2018; 33(2): 106-11.
12. Rochester L, Baker K, Hetherington V, Jones D, Willems AM, Kwakkel G et al. Evidence for motor learning in Parkinson's disease: acquisition, automaticity and retention of cued gait performance after training with external rhythmical cues. *Brain Res*. 2010; 1319: 103-11. doi: 10.1016/j.brainres.2010.01.001. Epub 2010 Jan 11.
13. Di Iorio W, Ciarimboli A, Ferriero G, Feleppa M, Baratto L, Matarazzo G et al. Action Observation

- in People with Parkinson's Disease. A Motor–Cognitive Combined Approach for Motor Rehabilitation. A Preliminary Report. *Diseases*. 2018; 6(3): 58. doi: 10.3390/diseases6030058.
14. Bek J, Holmes PS, Craig CE, Franklin ZC, Sullivan M, Webb J et al. Action Imagery and Observation in Neurorehabilitation for Parkinson's Disease (ACTION-PD): development of a user-informed home training intervention to improve functional hand movements. *Parkinson's Dis*. 2021; 2021: 4559519. doi: 10.1155/2021/4559519.
 15. Rabey JM, Korczyn AD. The Hoehn and Yahr Rating Scale for Parkinson's Disease. In: Przuntek, H., Kraus, P.H., Klotz, P., Korczyn, A.D. (eds) *Instrumental Methods and Scoring in Extrapyrmidal Disorders*. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-78914-4_2.
 16. de Morton NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother*. 2009; 55(2): 129-33. doi: 10.1016/s0004-9514(09)70043-1.
 17. Maher CG, Sherrington C, Herbert RD, Moseley AM, Elkins M. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther*. 2003; 83(8): 713-21.
 18. Braun S, Beurskens A, Kleynen M, Schols J, Wade D. Rehabilitation with mental practice has similar effects on mobility as rehabilitation with relaxation in people with Parkinson's disease: a multicentre randomised trial. *J Physiother*. 2011; 57(1): 27-34. doi: 10.1016/S1836-9553(11)70004-2.
 19. Tamir R, Dickstein R, Huberman M. Integration of motor imagery and physical practice in group treatment applied to subjects with Parkinson's disease. *Neurorehabil Neural Repair*. 2007; 21(1): 68-75. doi: 10.1177/1545968306292608.
 20. Santiago LMdM, de Oliveira DA, de Macêdo Ferreira LGL, de Brito Pinto HY, Spaniol AP, de Lucena Trigueiro LC et al. Immediate effects of adding mental practice to physical practice on the Gait of individuals with Parkinson's disease: randomized clinical trial. *NeuroRehabilitation*. 2015; 37(2): 263-71. doi: 10.3233/NRE-151259.
 21. Sarasso E, Agosta F, Piramide N, Gardoni A, Canu E, Leocadi M et al. Action Observation and Motor Imagery Improve Dual Task in Parkinson's Disease: A Clinical/fMRI Study. *Mov Disord*. 2021; 36(11): 2569-82.
 22. Lokhandwala M, Kulkarni V, Nair M. Comparison of Combined Effects of Physical and Mental Practice with Physical Practice alone on Functional Independence in Parkinson (PD) Patients. *Int J Physiother Res*. 2019; 7: 3207-13. doi: 10.16965/ijpr.2019.160.
 23. Abraham A, Hart A, Andrade I, Hackney ME. Dynamic neuro-cognitive imagery improves mental imagery ability, disease severity, and motor and cognitive functions in people with Parkinson's disease. *Neural Plast*. 2018; 2018: 6168507. doi: 10.1155/2018/6168507.
 24. Bovend'Eerdt TJ, Dawes H, Sackley C, Wade DT. Practical research-based guidance for motor imagery practice in neurorehabilitation. *Disabil Rehabil*. 2012; 34(25): 2192-2200. doi: 10.3109/09638288.2012.676703. Epub 2012 Apr 25.
 25. Nicholson V, Watts N, Chani Y, Keogh JW. Motor imagery training improves balance and mobility outcomes in older adults: a systematic review. *J Physiother*. 2019; 65(4): 200-207.
 26. Jackson PL, Lafleur MF, Malouin F, Richards C, Doyon J. Potential role of mental practice using motor imagery in neurologic rehabilitation. *Arch Phys Med Rehabil*. 2001; 82(8): 1133-41. doi: 10.1053/apmr.2001.24286.
 27. Oostra KM. Mental Practice through motor imagery in gait rehabilitation following acquired brain injury. Thesis: Ghent University; 2016.
 28. Gowda AS, Memon AN, Bidika E, Salib M, Rallabhandi B, Fayyaz H. Investigating the Viability of Motor Imagery as a Physical Rehabilitation Treatment for Patients With Stroke-Induced Motor Cortical Damage. *Cureus*. 2021; 13(3): e14001. doi: 10.7759/cureus.14001.
 29. Anil K, Hall SD, Demain S, Freeman JA, Ganis G, Marsden J. A Systematic Review of Neurofeedback for the Management of Motor Symptoms in Parkinson's Disease. *Brain Sci*. 2021; 11(10): 1292.
 30. Calvo A, Chiò A, Pagani M, Cammarosano S, Dematteis F, Moglia C et al. Parkinsonian traits in amyotrophic lateral sclerosis (ALS): a prospective population-based study. *J Neurol*. 2019; 266(7): 1633-42. doi: 10.1007/s00415-019-09305-0. Epub 2019 Apr 4.
 31. Abidi M, De Marco G, Grami F, Termoz N, Couillandre A, Querin G et al. Neural correlates of motor imagery of Gait in amyotrophic lateral sclerosis. *J Magn Reson Imaging*. 2021; 53(1): 223-33. doi: 10.1002/jmri.27335. Epub 2020 Sep 7.
 32. Bae YH, Ko Y, Ha H, Ahn SY, Lee W, Lee SM. An efficacy study on improving balance and Gait in subacute stroke patients by balance training with additional motor imagery: a pilot study. *J Phys Ther Sci*. 2015; 27(10): 3245-8. doi: 10.1589/jpts.27.3245. Epub 2015 Oct 30.
 33. Li RQ, Li ZM, Tan JY, Chen GL, Lin WY. Effects of motor imagery on walking function and balance in patients after stroke: A quantitative synthesis of randomized controlled trials. *Complement Ther Clin Pract*. 2017; 28: 75-84. doi: 10.1016/j.ctcp.2017.05.009. Epub 2017 May 26.
 34. Woodrow-Hill C, Poliakoff E, Gowen E, Vogt S, Sullivan M. Can action observation and/or motor imagery be used to improve computer-based actions in people with Parkinson's? In BNA 2021

- Festival of Neuroscience Poster abstracts. Vol. 5. Sage Publications Ltd. 2021. p. 235-236. (Brain and Neuroscience Advances). doi: 10.1177/23982128211035062.
35. Gil-Bermejo-Bernardez-Zerpa A, Moral-Munoz JA, Lucena-Anton D, Luque-Moreno C. Effectiveness of motor imagery on motor recovery in patients with multiple sclerosis: systematic review. *Int J Environ Res Public Health*. 2021; 18(2): 498. doi: 10.3390/ijerph18020498.
 36. Moshref-Razavi S, Sohrabi M, Sotoodeh MS. Effect of neurofeedback interactions and mental imagery on the elderly's balance. *Iran J Ageing*. 2017; 12(3): 288-99. doi: 10.21859/sija.12.3.288.
 37. Mirelman A, Rochester L, Maidan I, Del Din S, Alcock L, Nieuwhof F et al. Addition of a non-immersive virtual reality component to treadmill training to reduce fall risk in older adults (V-TIME): a randomised controlled trial. *Lancet*. 2016; 388(10050): 1170-82. doi: 10.1016/S0140-6736(16)31325-3. Epub 2016 Aug 11.
 38. Avanzino L, Pelosin E, Martino D, Abbruzzese G. Motor timing deficits in sequential movements in Parkinson disease are related to action planning: a motor imagery study. *PloS One*. 2013; 8(9): e75454. doi: 10.1371/journal.pone.0075454.
 39. Caligiore D, Mustile M, Spalletta G, Baldassarre G. Action observation and motor imagery for Rehabilitation in Parkinson's disease: A systematic review and an integrative hypothesis. *Neurosci Biobehav Rev*. 2017; 72: 210-22. doi: 10.1016/j.neubiorev.2016.11.005. Epub 2016 Nov 16.
 40. Taube W, Mouthon M, Leukel C, Hoogewoud HM, Annoni JM, Keller M. Brain activity during observation and motor imagery of different balance tasks: an fMRI study. *Cortex*. 2015; 64: 102-14. doi: 10.1016/j.cortex.2014.09.022. Epub 2014 Oct 27.
 41. Kim JY, Kim JM, Ko EY. The effect of the action observation physical training on the upper extremity function in children with cerebral palsy. *J Exerc Rehabil*. 2014; 10(3): 176-183. doi: 10.12965/jer.140114.
 42. Sarasso E, Agosta F, Chiesi M, Piramide N, Canu E, Ravani I et al. Clinical and fMRI Effects of Action Observation and Motor Imagery Training on Dual-Task Performances in Parkinson's Disease Patients With Postural Instability and Gait Disorders (P5. 6-010). *Neurology*. 2019; 92(Suppl 15): P5.010.

