

Incidence of Fodder Cutter (Toka) Hand Injury and their Management in D.I. Khan, Pakistan

Najeebullah¹, Muhammad Shafiq^{1*}, Syed Wasif Ali Naqvi¹, Muhammad Hamyun², Muhammad Ali³, Shahjahan⁴

ABSTRACT

OBJECTIVE: This study aimed to determine the incidence of toka hand injuries by age, gender, and occupation and their management in D.I. Khan.

METHODOLOGY: This descriptive study was conducted from January to December 2023 in the Department of Orthopaedics, DHQ Teaching Hospital, D.I. Khan, Pakistan. Sixty-six patients presenting to the hospital during 2023 with toka hand injuries were selected. Demographic variables were age, gender, and occupation. Research variables were the site of injury and mode of management. All data was analyzed using SPSS version 29.

RESULTS: Out of 66 patients, Toka injuries were predominant in females 39(59.09%). Most patients, 51 (77.27%), were aged below 30 years, and 42(63.64%) were non-professional by occupation. The most common site of injury was distal to the wrist 33(50%), while it was 27(40.91%) at the wrist level, and 6 (9.09%) were proximal to the wrist. Debridement and stump formation was done in 40(60.60%) while 26 (39.40%) patients underwent fixation and stabilization.

CONCLUSION: Toka machine injuries mainly result in hand amputations. Safety features must be identified in this region, workers must be adequately educated, and work-related ergonomics must be ensured. Further advancements in management are also needed.

KEYWORDS: Toka injuries, Hand injuries, D. I. Khan, Fodder cutter, Stump, Amputation, Agriculture

INTRODUCTION

Globally, occupational injuries are responsible for about 10.5 million Disability-adjusted life years (DALYs)¹. The International Labor Organization (ILO) estimates that there are around 340 million occupational accidents worldwide². Hand injuries are the leading cause of lifelong disability in agriculture-related trauma worldwide³. One standard machine used in agriculture is the chaff cutter, which can cause permanent injuries to the upper limb (hand)⁴. A study in India on agricultural-related injuries reported that 45% of injuries were due to chaff-cutter incidents⁵. Toka or Fodder (chaff) cutter hand injuries are common in rural areas of Pakistan due to farmers use of fodder cutter machines for food preparation for their cattle⁶. Farmers cut hay into small pieces by using Toka machines. Fingers get trapped into the cutting wheel of the roller, which will result in the crushing or cutting of fingers and hands between the rollers⁷.

¹Department of Orthopedics, Gomal Medical College, D.I. Khan, KPK-Pakistan

²Department of Surgery, Gomal Medical College, D.I. Khan, KPK-Pakistan

³Department of Forensic, Gomal Medical College, D.I. Khan, KPK-Pakistan

⁴Department of Biochemistry, Gomal Medical College, D.I. Khan, KPK-Pakistan

Correspondence: drshafiqorthosurg@gmail.com
doi: 10.22442/jlumhs.2024.01182

Received: 16-08-2024

Revised: 24-09-2024

Accepted: 24-09-2024

Published Online: 26-09-2024

Each piece of machinery produces a spectrum of soft-tissue injuries.

Bone damage is characterized by the loss of a digit or limb, loss of function with permanent disability and serious complications⁸. A survey conducted from November 2022 to April 2023 involved 55 respondents, predominantly from Sindh and Punjab, with findings indicating 507 agricultural injuries where fodder cutter with 201 injuries was a primary source accounting for approximately 40% of all injuries⁹. A study in Sialkot by Mehmood R 2015¹⁰ showed that the toka machine was primarily involved in agricultural equipment found in 17 (51.5%) out of 33 patients with agriculture-related injuries. People of all ages and genders are affected by toka machines¹¹. While playing with the toka machine, the children get trapped in their fingers and hands. The main aim of reducing this risk is a quick and precise initial assessment. Injury becomes mismanaged when non-specialists need help understanding the peculiarity of the injuries methodically, initially in the emergency¹². No significant data is available on different search engines for Toka injuries in D.I. Khan. For this reason, the study was conducted to determine the incidence of toka hand injuries in age, gender, and occupation, along with their management in D. I. Khan. The objectives of this study were to determine the incidence of fodder cutter (Toka) hand injuries based on age, gender, and occupation, along with their management in D.I. Khan, Pakistan.





METHODOLOGY

This descriptive cross-sectional study was conducted from January to December 2023 in the Department of Orthopedics, DHQ Teaching Hospital, D.I. Khan, Pakistan. Sixty-six patients presenting to the hospital with a history of toka hand injuries were selected. A consecutive sampling technique was used. Patients presenting to the hospital with toka hand injuries were included, and all other patients were excluded. Data of all toka injury patients from January to December 2023 was retrieved from hospital records. Demographic variables were age in two groups (below 30 years and above 30 years), gender in two groups (male and female) and occupation in two groups (professional and non-professional). Research variables were the injury site in three groups (distal to the wrist, at wrist level and proximal to the wrist) and the management mode in two groups (debridement & stump formation and fixation & stabilization). Count and percentage described sample distribution. Population distribution was represented by 95% CI using the standard approximation method through the online statistical calculator¹³. The comparison was based on overlapping with confidence intervals. All data was analyzed using SPSS version 29.

RESULTS

Descriptive Analysis and Estimation of Parameters

Distribution of Toka injury by age, gender, occupation, site of injury and mode of management: Out of 66 patients with toka injury, 51(77.27%) were aged below

30 years & 15(22.73%) were over 30 years. Twenty-seven (40.91%) were males & 39(59.09%) were females. Twenty-four (36.36%) were professional & 42 (63.64%) were non-professional. Thirty-three (50%) were distal to the wrist, 27(40.91%) at the wrist level & 6(9.09%) were proximal to the wrist. Forty patients (60.60%) underwent debridement & stump formation, while 26(39.40%) underwent fixation & stabilization.

In our study, toka injury was higher in females (59.09%, 95% CI 47.23-70.95) than males (40.91%, 95% CI 29.09-52.77), higher in the age group below 30 years (77.27%, 95% CI 67.16-87.38) than above 30 years (22.73%, 95% CI 12.62-32.84), higher in non-professional (63.33%, 95% CI 52.03-75.24) than professional (33.33%, 95% CI 24.76-47.96), highest at level distal to wrist 33(50%, 95% CI 37.94-62.06) than at wrist 27(40.91%, 95% CI 29.05-52.77) & proximal to wrist 6(9.09%, 95% CI 2.15-16.03). Most of the patients undergone debridement & stump formation 40(60.60%, 95% CI 48.82-72.39), while 26(39.40%, 95% CI 27.61-51.18) patients undergone fixation & stabilization as shown in **Table I**.

DISCUSSION

Distribution of Toka injury by age groups:

In our study, toka injury was more prevalent in age groups below 30 years 51(77.27%, 95% CI 67.16-87.38) than above 30 years 15(22.73%, 95% CI 12.62-32.84). Similar to our study, higher prevalence in the age group below 30 years was reported by Choudhry NA 2016¹³ from southern Punjab out of 73 patients of

TABLE I: DISTRIBUTION OF TOKA INJURY BY AGE, GENDER, OCCUPATION, SITE OF INJURY AND MODE OF MANAGEMENT IN THE POPULATION OF D. I. KHAN, PAKISTAN (n=66)

Variables	Attributes	Sample Statistics		95% CI for proportion	
		Counts	Percentage	Lower	Upper
Age	Below 30 years	51	77.27%	67.16%	87.38%
	Above 30 years	15	22.73%	12.62%	32.84%
Gender	Male	27	40.91%	29.05%	52.77%
	Female	39	59.09%	47.23%	70.95%
Occupation	Professional	24	36.36%	24.76%	47.97%
	Non- professional	42	63.64%	52.03%	75.24%
Site of Injury	Distal to wrist	33	50%	37.94%	62.06%
	At wrist level	27	40.91%	29.05%	52.77%
	Proximal to wrist	6	9.09%	2.15%	16.03%
Mode of Management	Debridement & stump formation	40	60.61%	48.82%	72.39%
	Fixation and stabilization	26	39.39%	27.61%	51.18%

toka injury, including 56(76.71%, 95% CI 67.02-86.41) below the age of 30 years & 17(23.29%, 95% CI 13.59-32.98) patients 31-60 years, and by Tunio ZH et al.¹⁴ from Sindh out of 56 patients affected by fodder cutter injury, where 36(64.29%, 95% CI 51.74-76.84) were aged below 30 years & 20(35.71%, 95% CI 23.16-48.26) were aged 31-60 years.

Unlike our study, a study by Ahmad et al.¹⁶ from Sialkot, out of 196 patients, showed lower prevalence in the age group below 30 years as 96(48.97%, 95% CI 41.98-55.98) and higher prevalence in the 31-60 year age group as 100(51.02%, 95% CI 44.02-58.02).

Distribution of Toka injury by gender

In our study, toka injury was higher in females 39 (59.09%, 95% CI 47.23-70.95) than males 27(40.91%, 95% CI 29.09-52.77). Similar to our study, a higher prevalence in females was reported by Choudhry NA 2016¹⁴, including 41(56.16%, 95% CI 44.78-67.55) females and 32(43.84%, 95% CI 32.45-55.25) males out of 73 patients.

Unlike our study, higher prevalence in male and lower in females was reported by Wanjara S 2022¹⁶ from Kenya, where 46(98%, 95% CI 93.75-100) were males & 1(2%, 95% CI 0.01-6.25) female out of 47 patients, by Tunio ZH et al.¹⁴ where 38(67.86%, 95% CI 55.63-80.09) were males & 18(32.14%, 95% CI 19.91-44.37) were females out of 56 patients, by Ahmad S 2018¹⁵ where 129(65.82%, 95% CI 59.18-72.46) were males & 67(34.18%, 95% CI 27.54-40.82) were females, by Ali D 2021¹⁷ from Lahore out of 30 cases of toka injury where 23(76.6%, 95% CI 61.53-91.80) males & 07 (23.4%, 95% CI 8.19-38.47) females, and by Bhatti SH 2021¹⁸ from Sindh out of 193 patients where 133 (68.91%, 95% CI 62.38-75.44) were male and 60(31.08%, 95% CI 24.56-37.62) female.

Distribution of Toka injury by occupation

In our study, the incidence of injury was higher in non-professional patients (63.33%, 95% CI 52.03-75.24) than in professionals (33.33%, 95% CI 24.76-47.96). No study showing Toka injury by occupation could be found. The non-professionals are casual workers unfamiliar with these machines, which may cause high incidence.

Distribution of Toka injury by site of injury

In our study, toka injury was highest at the level distal to wrist 33(50%, 95% CI 37.94-62.06) than at wrist 27 (40.91%, 95% CI 29.05-52.77) & proximal to wrist 6 (9.09%, 95% CI 2.15-16.03). Our findings were somewhat similar to the study by Choudhry NA 2016¹⁴, with 24(32.88%, 95% CI 22.10-43.65) distal to the wrist, 22(30.14%, 95% CI 19.61-40.66) at the wrist while the highest prevalence at the level proximal to the wrist in 27(36.98%, 95% CI 25.91-48.06) patients. Also, our findings were somewhat similar to the study by Bhatti SH 2021¹⁸ with 66 (34.20%, 95% CI 27.50-540.89) distal to the wrist and 34 (17.62%, 95% CI 12.24-22.99) proximal to the wrist. However, the highest prevalence was at wrist 94 (48.18%, 95% CI 41.65-55.76).

Distribution of Toka injury by mode of management

Toka injuries are a major contributor to disability in the agriculture sector of Pakistan. In our study, most of the patients undergone debridement & stump formation 40(60.60%, 95% CI 48.82-72.39), while 26 (39.40%, 95% CI 27.61-51.18) undergone fixation and stabilization. Unlike our study, Choudhry NA 2016¹⁴ reported higher debridement & stump formation in 70 (96%, 95% CI 91.34-100), and only 3(4%, 95% CI 0.44-8.66) patients underwent fixation & stabilization. Similar to our study, Wanjara S 2022¹⁶ reported 33 (70.21%, 95% CI 57.14-83.29) debridement & stump formation and 14(29.79%, 95% CI 16.71-42.86) fixation & stabilization out of 47 patients, Ali D 2021¹⁷ reported 18(60.1%, 95% CI 42.47-77.53) debridement & stump formation and 12(39.9%, 95% CI 22.47-57.53) fixation & stabilization out of 30 patients of toka injury, and Tunio ZH et al.¹⁴ reported 34(60.71%, 95% CI 47.92-73.51) debridement & stump formation and 22(39.29%, 95% CI 26.49-52.08) fixation & stabilization out of 56 patients.

CONCLUSION

Toka or fodder cutter machine injuries are common in our settings. Most of that results in amputations and permanent disabilities, suggesting increased dependency. Non-professional and non-expert persons of a young age are commonly involved in these injuries.

RECOMMENDATIONS

Awareness should be started in print, electronic, and social media. Safety/protective features such as power cut-off, limit switch, trigger/belt deflector system, front shield grill, belt cover, gear cover, wooden foundation, and protection barrier plates must be identified. 20 Proper education of workers in this region, work-related ergonomics, and training of paramedical staff working in the peripheral hospitals to handle these injuries properly are also needed.

Ethical permission: Gomal Medical College, D. I. Khan, Khyber Pakhtunkhwa, Pakistan, ERC letter No. 225/GJMC/JC.

Conflict of Interest: No conflicts of interest, as stated by authors.

Financial Disclosure / Grant Approval: No funding agency was involved in this research.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION

All authors contributed equally to the study.

REFERENCES

1. Concha-Barrientos M, Nelson DI, Fingerhut M, Driscoll T, Leigh J. The global burden due to occupational injury. *Am J Ind Med.* 2005; 48(6):

- 470-481. doi: 10.1002/ajim.20226.
2. International Labour Organization (ILO). World Statistics. Available from: <https://www.ilo.org/moscow/areas-of-work/occupational-safety-and-health/WCMS>.
 3. Mucci N, Traversini V, Lulli LG, Baldassarre A, Galea RP, Arcangeli G. Upper Limb's Injuries in Agriculture: A Systematic Review. *Int J Environ Res Public Health*. 2020; 17(12): 4501. doi: 10.3390/ijerph17124501.
 4. Asha KR, Kumar A, Singh JK, Kushwaha HL. Sensor-based Safety Alarm System for Injury Prevention in Chaff cutter machine. *Smart Agri Technol*. 2023; 5(1): 100282. doi: 10.1016/j.atech.2023.100282.
 5. Chopra S, Kumar P, Kumar M. Severity of agricultural injuries and its economic consequences in the border belt of Gurdaspur District of Punjab, India. *J Agromedicine*. 2024; 29(2): 179-188. doi: 10.1080/1059924X.2023.2281517. Epub 2023 Nov 16.
 6. Wikipedia. Chaff Cutter. Wikipedia. Available from: https://en.wikipedia.org/wiki/Chaff_cutter.
 7. Shrestha KM, Shrestha B, Kandel PR, Baral R, Pandey A, Singh GP. Fodder Cutter Machine Injuries of Hand. *J Univers Coll Med Sci*. 2018; 6: 14-18. doi: 10.3126/jucms.v6i1.21693.
 8. Yaffe MA, Kaplan FT. Agriculture injuries to hand and upper extremity. *J Am Acad Orthop Surg*. 2014; 22(10): 605-613. doi: 10.5435/JAAOS-22-10-605.
 9. Raza MMS, Tunio ZH, Ujjan ID, Issa SF. Insights into Agricultural Machine Injuries in Pakistan: An Orthopedic Surgeons Survey (2022–2023). *Safety*. 2024; 10(3): 55. doi: 10.3390/safety10030055.
 10. Mehmood R, Aziz S, Jehan S, Qureshi MA. Agricultural related injuries; Spectrum & management outcome in General Surgical Unit. *Professional Med J*. 2015; 22(2): 175-180. doi: 10.29309/TPMJ/2015.22.02.1369.
 11. New-Aaron M, Semin J, Duysen EG, Madsen M, Musil K, Rautiainen RH. Comparison of agricultural injuries reported in the media and census of fatal occupational injuries. *J Agromedicine*. 2019; 24(3): 279-287. doi: 10.1080/1059924X.2019.1593276. Epub 2019 Mar 30.
 12. Statistics Kingdom. Proportion confidence interval calculator. Statistics Kingdom: Melbourne, Australia 2007. Available from: <https://www.statskingdom.com/proportion-confidence-interval-calculator.html>.
 13. Choudhry NA, Azeem M, Iqbal MZ, Choudhry AA. Fodder Cutter Amputations-Evaluation of Risk Factors. *J Pak Orthop Assoc*. 2016; 28(2): 46-49.
 14. Tunio ZH, Jhatiyal RA, Jokhio MF, Ali SM, Baloch RA, Makhdoom A. Fodder Cutter Machine Injuries: An Unsafe Household Item in Rural Sindh, Pakistan. *J Liaq Uni Med Health Sci*. 2021; 20(02): 96-99. doi: 10.22442/jlumhs.2021.00813.
 15. Ahmad S, Hamid K, Shabir F, Latif A. Pattern of Fodder Cutting Machine Injuries in Sialkot Region. *Pak J Med Health Sci*. 2018; 12(4): 1368-1370.
 16. Wanjara S, Oduor P. Hand injuries in Kenya: a chaff cutter menace. *F1000Res*. 2022; 11: 1255. doi: 10.12688/f1000research.126590.1.
 17. Ali D, Kaiser MA, Khokhar MI, Zeb A. Fodder cutter (Tokka) injuries: A preventable morbidity. *J Pak Med Assoc*. 2021; 71(3): 1022-1024.
 18. Bhatti SH, Abbasi MK, Naqvi ASAH. Pattern of fodder cutting machine injury in Gambat region, Sindh, Pakistan. *Rawal Med J*. 2021; 46(3): 560-563.
 19. Yaseen MU, Saddique G, Sabar MA, Ashraf M, Ahmed S, Ahmed M. Development and installation of safety features in fodder chopper to make its operation reliable and hazard free. *J Agric Res*. 2022; 60(1): 59-66.

