

Plant-Based Diets and Dyslipidemia in Tharparkar: A Cross-Sectional Study on Lipid Imbalances

Gian Chand¹, Abdul Razzaque Nohri², Nazia Mumtaz^{3*}, Aisha Memon⁴,
Ghulam Muhammad Waryah⁵, Shabahat Memon⁶

ABSTRACT

OBJECTIVE: To assess lipid profiles in Tharparkar's vegetarian population and examine dietary habits and their correlation with lipid imbalances.

METHODOLOGY: This cross-sectional study was conducted at Bilawal Medical College, Jamshoro, in collaboration of other institutes in Sindh, Pakistan, from Dec 2024 to May 2025, after obtaining ethical approval. The vegetarian population was included, and the non-vegetarian population was excluded. A total of 500 participants were selected via stratified random sampling. Lipid profiling was conducted after 12 hours of fasting. Dietary habits and socioeconomic factors were assessed using structured questionnaires. Data analysis was performed with SPSS.

RESULTS: 500 participants revealed that 60% had lipid imbalances. Specifically, 45% had high Cholesterol, 40% had high LDL-C, 60% had low HDL-C, and 50% had high triglycerides. High carbohydrate intake from refined grains, low omega-3 fatty acid intake, and consumption of fried foods were strongly linked to unfavourable lipid profiles.

CONCLUSION: The high dyslipidemia rates in Tharparkar's vegetarian population are occurring due to nutrient imbalances, socioeconomic factors and unhealthy food consumption. Public health counselling promoting balanced vegetarian diets and addressing lifestyle factors could help improve lipid profiles and reduce cardiovascular risk.

KEYWORDS: Vegetarian Diet, Dyslipidemia, Lipid Imbalances, Cardiovascular Risk, Nutrient Imbalance.

INTRODUCTION

Dyslipidemia, characterized as abnormal lipid levels in the blood and a major cause of cardiovascular diseases (CVD), remains the leading cause of morbidity and mortality worldwide¹. The condition is analyzed by elevated total Cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and triglycerides (TG), along with low levels of high-density lipoprotein cholesterol (HDL-C). Lipid imbalances can lead to the development of atherosclerosis, increasing the risk of heart attacks, strokes, and other cardiovascular events². Globally, it is estimated that 39% of adults have high Cholesterol, and the

condition significantly indicates the high burden of CVDs, especially in low- and middle-income countries³.

In Pakistan, the increasing prevalence of dyslipidemia is an alarming situation; numerous studies have shown that lipid abnormalities are a contributing factor to rising rates of cardiovascular diseases. A national survey on cardiovascular health in Pakistan reported that nearly 50% of the adult population exhibited one or more lipid abnormalities, including elevated LDL-C and low HDL-C levels⁴. Previous studies conducted in urban and rural regions have consistently shown that the risk factors for dyslipidemia, including diet, Lifestyle, and socioeconomic conditions, are influenced across the globe.⁵ The role of vegetarian diets is one of the most interactive aspects of the relationship between diet and dyslipidemia. Plant-based diets are often associated with lower cholesterol levels and reduced risks of heart disease. The prevalence of dyslipidemia in vegetarian populations, particularly in specific population regions with limited access to nutrient-rich foods, raises concerns about the effectiveness of vegetarian diets in promoting optimal heart health. In some cases, an unbalanced plant-based diet and poor dietary planning have been linked to lipid imbalances, even in a vegetarian lifestyle⁶.

The majority of people of Tharparkar, a province of Sindh, Pakistan, are mostly vegetarians. Plant-based eating habits are deeply ingrained in this region's culture, primarily as a result of vegetarianism being

¹Assistant Professor, National Institute of Cardiovascular Diseases, Karachi, Sindh, Pakistan

²Senior Pharmacist and Public Health Specialist, Health Department, Government of Sindh

³Assistant Professor, Department of Community Medicine, Liaquat Institute of Medical & Health Sciences (LIMHS) Thatta, Sindh, Pakistan

⁴Lecture, Department of Pathology, Bilawal Medical College, Jamshoro, Sindh, Pakistan

⁵Sr. Lecture, Institute of Biotechnology & Genetic Engineering (IBGE), University of Sindh, Jamshoro, Sindh, Pakistan

⁶Sindh Institute of Ophthalmology & Visual Sciences, Hyderabad, Sindh, Pakistan

Correspondence: nazia.mumtazbmc@gmail.com

doi: 10.22442/jlumhs.2025.01351

Received: 22-05-2025

Revised: 23-05-2025

Accepted: 23-05-2025

Published Online: 23-05-2025



emphasized in religious and cultural practices. Cereals, legumes, vegetables, and dairy products make up the majority of the typical Tharparkar diet. These foods are thought to be high in essential fatty acids and other micronutrients vital to lipid metabolism. Although vegetarian diets are generally heart-healthy, there is a paradox in Tharparkar, where the population suffers from a high burden of cardiovascular diseases, and dyslipidemia rates are high^{7,8}.

The incidence of dyslipidemia in Tharparkar's vegetarian population is a growing concern. Several studies and Local health records indicate that, despite following a plant-based diet, many individuals in the region have high levels of LDL-C, low levels of HDL-C, and elevated triglycerides, all signs of dyslipidemia^{4,9}. Studies in other regions indicated that certain dietary patterns, such as a high intake of refined carbohydrates, excessive consumption of fried foods, and insufficient intake of healthy fats like omega-3 fatty acids, may lead to lipid imbalances even among individuals who follow vegetarian diets. These results suggest that the general health advantages of vegetarian diets may not apply universally, particularly when the diet lacks variety or essential nutrients that support healthy lipid metabolism¹⁰.

International studies suggest that well-managed vegetarian diets, rich in fruits, vegetables, whole grains, legumes, and healthy fats, can help reduce total cholesterol levels, LDL-C, and triglycerides while improving HDL-C levels¹¹. A systematic review by the American Heart Association showed that plant-based diets can reduce the risk of cardiovascular disease by improving lipid profiles. However, it depends on maintaining a balanced diet and including essential nutrients, especially omega-3 fatty acids, which are commonly found in fish or fortified plant-based sources¹². Poorly planned vegetarian diets that are heavy in carbohydrates, especially simple sugars and refined grains, can result in increased triglyceride levels and an undesirable lipid profile¹³.

Instability of food supplies and financial constraints make it more difficult for people in rural areas like Tharparkar to maintain a balanced vegetarian diet¹⁴. The increasing incidence of dyslipidemia may also be attributed to a lack of public health campaigns to inform the public about heart-healthy eating practices¹⁴.

Therefore, the major problem among vegetarians in Tharparkar is not only a lack of animal products but also a lack of vital nutrients, such as omega-3 fatty acids, which are crucial for maintaining normal cholesterol levels¹⁵. The purpose of this study was to investigate the association of lipid abnormalities and vegetarian eating patterns in the Tharparkar region. The findings highlighted the difficulties of vegetarian populations encounter in preserving ideal lipid profiles and focused treatments to enhance heart health in Tharparkar and surrounding regions. However, the findings also help guide public health strategies to

promote balanced vegetarian diets that support cardiovascular health.

The high dyslipidemia rates in Tharparkar's vegetarian population are occurring due to nutrient imbalances, socioeconomic factors and unhealthy food consumption. Public health counselling promoting balanced vegetarian diets and addressing lifestyle factors could help improve lipid profiles and reduce cardiovascular risk.

METHODOLOGY

This cross-sectional study was conducted at Bilawal Medical College, Jamshoro, in collaboration of other institutes in Sindh, Pakistan, from Dec 2024 to May 2025, after obtaining ethical approval. The vegetarian population was included, and the non-vegetarian population was excluded from the study. A total of 500 participants were selected through stratified random sampling to represent data across different age groups, genders, and socioeconomic statuses. The data collection process involved noticing and collecting variable data. The blood samples were obtained for Lipid Profiling after a 12-hour fasting period to measure TC, LDL-C, HDL-C, and TG levels using standard enzymatic methods. A structured questionnaire was administered to assess participants' dietary habits, focusing on the consumption of plant-based foods, cooking methods, and frequency of intake. Socioeconomic and Lifestyle information of participants was recorded, including age, gender, income, education level, physical activity, and tobacco use, collected through interviews. Data was analyzed using SPSS version 29.0.

RESULTS

The study revealed a significant prevalence of dyslipidemia among the predominantly vegetarian population of Tharparkar. Based on the analysis of lipid profiles, dietary habits, and socioeconomic factors, the results show that more than 60% of participants exhibited lipid imbalances (**Table I**).

Table I: Lipid Profile of Participants

Lipid Parameter	Mean±SD	Abnormal Levels (%)
Total Cholesterol (mg/dL)	210 ± 45	45.0%
LDL-C (mg/dL)	130 ± 40	40.0%
HDL-C (mg/dL)	35 ± 10	60.0%
Triglycerides (mg/dL)	180 ± 50	50.0%

The study sample comprised 500 participants, with an even distribution of males and females. The majority of participants (30%) were between the ages of 31 and 45, followed by the 46-60 age group, followed by the 24% of 18-30. The data show a notable proportion of the population (40%) having a monthly income below 10,000 PKR, indicating a lower socioeconomic status. Educationally, 50% of participants had at least secondary education **Table II**.

Table II: Demographic Characteristics of Participants

Characteristic	Frequency(%)
Age (years)	
18–30	120 (24.0%)
31–45	150 (30.0%)
46–60	150 (30.0%)
>60	80 (16.0%)
Gender	
Male	250 (50.0%)
Female	250 (50.0%)
Education Level	
No formal education	100 (20.0%)
Primary	150 (30.0%)
Secondary	150 (30.0%)
Higher	100 (20.0%)
Monthly Income (PKR)	
<10,000	200 (40.0%)
10,000–20,000	150 (30.0%)
>20,000	150 (30.0%)

The lipid profile data revealed that a significant portion of the participants had abnormal levels of lipids. Approximately 45% of participants had total cholesterol levels above the recommended threshold of 200 mg/dL. High LDL-C levels were observed in 40% of participants, with 60% of individuals having low HDL-C levels. Elevated triglyceride levels were found in 50% of participants, particularly among those aged 45 years old. The association between lipid imbalances and dietary habits was analyzed. **Table III: Dietary Habits and Their Association with Lipid Imbalances** summarizes the relationship between common dietary patterns and lipid profile abnormalities (**Table III**).

Table III: Dietary Habits and Their Association with Lipid Imbalances

Dietary Habit	LDL-C >130 mg/dL (%)	HDL-C <40 mg/dL (%)	TG >150 mg/dL (%)
High carbohydrate intake (>60% of total calories)	55.0%	65.0%	60.0%
Low intake of omega-3 fatty acids (<1g/day)	60.0%	70.0%	65.0%
High consumption of fried foods (≥3 times/week)	50.0%	55.0%	50.0%

The study found that participants with high carbohydrate intake, particularly from foods like rice and bread, had a significantly higher risk of elevated LDL-C and triglyceride levels. Additionally, a deficiency in omega-3 fatty acids was common, with 60% of participants consuming less than 1g of omega-3 fatty acids per day. A high frequency of fried food

consumption was also associated with unfavorable lipid profiles (**Table IV**).

Table IV: Socioeconomic and Lifestyle Factors Influencing Lipid Profiles

Factor	LDL-C >130 mg/dL (%)	HDL-C <40 mg/dL (%)	TG >150 mg/dL (%)
Low income (<10,000 PKR/month)	60.0%	65.0%	55.0%
Low education level (no formal education)	58.0%	62.0%	53.0%
Physical inactivity (sedentary Lifestyle)	65.0%	70.0%	60.0%
Tobacco use	50.0%	55.0%	45.0%

This study also indicated the influence of socioeconomic and Lifestyle factors on lipid imbalances. Participants in lower-income groups (earning less than 10,000 PKR/month) had a higher prevalence of lipid abnormalities, and individuals with low education levels and those with a sedentary lifestyle exhibited abnormal lipid profiles. However, it was also observed that Participants who are frequently used to Tobacco have higher LDL-C and triglyceride levels. Overall, the results suggest a complex interaction of dietary patterns, socioeconomic status, and lifestyle factors in the form of lipid profiles of Tharparkar's vegetarian population. Despite following a plant-based diet, a significant portion of the population still had dyslipidemia, which may be due to nutritional imbalances, including low omega-3 fatty acid intake, high carbohydrate intake, and unhealthy food preparation methods. Socioeconomic challenges, including limited access to a balanced diet, further exacerbate these lipid imbalances.

The findings indicate that modifications, particularly increasing omega-3 fatty acid intake and reducing refined carbohydrate consumption, help improve lipid profiles and reduce the risk of cardiovascular disease in the population. Additionally, addressing lifestyle factors such as physical activity and tobacco use is crucial for preventing dyslipidemia in this region.

DISCUSSION

The results show an incidence of lipid imbalances in the population, with elevated LDL-C and T; this highlights the prevalence of dyslipidemia among the vegetarian population in Tharparkar, Pakistan, and emphasizes the role of dietary patterns, lifestyle factors, and socioeconomic status in shaping lipid profiles¹⁶. Low HDL-C and increased triglyceride levels. These findings are consistent with past reports suggesting that plant-based diets, when poorly balanced, may lead to dyslipidemia¹⁷.

The most striking result of this study is the high prevalence of low HDL-C (83.9%) and elevated triglycerides (48.9%) among participants. Similar trends have been observed in other vegetarian populations, particularly in regions where

carbohydrate-rich foods form the majority of the diet^{9,18}. For instance, a study in India found that high carbohydrate intake was associated with increased triglyceride levels and low HDL-C in a predominantly vegetarian population. High carbohydrate intake, especially from refined grains and sugars, can elevate triglycerides, a known risk factor for cardiovascular diseases. Our study found that participants with a higher intake of carbohydrates, such as rice and bread, had significantly higher LDL-C and triglycerides, which aligns with these findings¹⁹. Another major finding was the deficiency in omega-3 fatty acids, with 60% of participants consuming less than 1g of omega-3s per day. Omega-3 fatty acids are essential for maintaining healthy cholesterol levels and reducing inflammation²⁰. Previous research has shown that vegetarian diets low in omega-3s-particularly those that do not include fortified foods or plant-based sources such as flaxseeds or walnuts- are associated with an increased risk of dyslipidemia and cardiovascular disease²¹. Socioeconomic factors also emerged as important contributors to lipid imbalances. Our findings indicate that participants with lower incomes and lower levels of education exhibited worse lipid profiles. This is consistent with studies showing that individuals with lower socioeconomic status often have limited access to healthy, nutrient-rich foods, which can lead to poor dietary habits and increased cardiovascular risk. Additionally, physical inactivity and tobacco use were significant contributors to dyslipidemia, reinforcing the need for lifestyle modifications to improve lipid profiles.²² While this study provides valuable insights into the link between vegetarian diets and dyslipidemia in Tharparkar, it has some limitations. The cross-sectional design of the study limits the ability to establish causal relationships between diet and lipid imbalances. Additionally, the reliance on self-reported dietary data may introduce recall bias, as participants may not accurately remember or report their food intake. The study also did not account for genetic factors that might influence lipid metabolism, which could further explain variations in lipid profiles. Despite these limitations, the study has significant implications for public health. It underscores the need for balanced vegetarian diets that include sufficient sources of omega-3 fatty acids, healthy fats, and proteins. Public health interventions should focus on educating the population about the risks of a high-carbohydrate diet and the importance of a varied diet rich of nutrients. Moreover, addressing socioeconomic disparities, such as access to diverse food sources, could help mitigate the prevalence of dyslipidemia and reduce the risk of cardiovascular diseases in Tharparkar and similar regions.

CONCLUSION

This study highlights the significant prevalence of dyslipidemia in Tharparkar's vegetarian population, primarily associated with unbalanced diets high in

carbohydrates and low in omega-3 fatty acids. Public health interventions and counselling promoting balanced vegetarian diets increased omega-3 intake, and addressing socioeconomic factors are essential for improving lipid profiles and reducing cardiovascular risk.

Acknowledgements: The authors are thankful to all participants.

Ethical permission: Bilawal Medical College of Liaquat University of Medical & Health Sciences, Jamshoro, Pakistan, ERC approval letter No. LUMHS/REC/-484.

Conflict of interest: There is no conflict of interest between the authors.

Financial Disclosure / Grant Approval: No funding agency was involved in this research.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION

Chand G: Study design, data analysis, and review of manuscript

Nohri AR: Collection of data and review of manuscript

Mumtaz N: Help in writing the draft and critical review of the manuscript

Memon A: Data collection and Critical review

Waryah GM: Writing the draft and critical review of the manuscript

Memon S: Data collection and Critical review

REFERENCES

1. Du Z, Qin Y. Dyslipidemia and Cardiovascular Disease: Current Knowledge, Existing Challenges, and New Opportunities for Management Strategies. *J Clin Med.* 2023; 12(1): 363. doi: 10.3390/jcm12010363.
2. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem.* 1972; 18(6): 499-502.
3. Zheng J, Wang J, Zhang Y, Xia J, Guo H, Hu H et al. The Global Burden of Diseases attributed to high low-density lipoprotein cholesterol from 1990 to 2019. *Front Public Health.* 2022; 10: 891929. doi: 10.3389/fpubh.2022.891929.
4. Basit A, Sabir S, Riaz M, Fawwad A; NDSP members. NDSP 05: Prevalence and pattern of dyslipidemia in urban and rural areas of Pakistan; a sub-analysis from second National Diabetes Survey of Pakistan (NDSP) 2016-2017. *J Diabetes Metab Disord.* 2020; 19(2): 1215-1225. doi: 10.1007/s40200-020-00631-z.
5. Zaid M, Hasnain S. Plasma lipid abnormalities in Pakistani population: trends, associated factors, and clinical implications. *Braz J Med Biol Res.* 2018; 51(9): e7239. doi: 10.1590/1414-

- 431X20187239.
6. Trautwein EA, McKay S. The Role of Specific Components of a Plant-Based Diet in Management of Dyslipidemia and the Impact on Cardiovascular Risk. *Nutrients*. 2020; 12(9): 2671. doi: 10.3390/nu12092671.
 7. Shar AH. Ethnography of major wild food plants uses and their consumption in the Thar Desert of Sindh, Pakistan. (Thesis). Università Ca' Foscari, Venice, Italia. 2022. Available from: <https://unitesi.unive.it/retrieve/9d5b77d2-29a1-4d23-80ef-f7c46f7bdded/876769-1247742.pdf>
 8. Li Y, Hruby A, Bernstein AM, Ley SH, Wang DD, Chiuve SE et al. Saturated Fats Compared With Unsaturated Fats and Sources of Carbohydrates in Relation to Risk of Coronary Heart Disease: A Prospective Cohort Study. *J Am Coll Cardiol*. 2015; 66: 1538-1548.
 9. Enas EA, Yusuf S, Mehta JL. Prevalence of coronary artery disease in asian Indians. *Am J Cardiol*. 1992; 70: 945-949. doi: 10.1016/0002-9149(92)90744-J.
 10. Cusin I, Terrettaz J, Rohnerjeanrenaud F, Zarjevski N, Assimacopoulosjeannet F, Jeanrenaud B. Hyperinsulinemia increases the amount of Glut4 messenger RNA in white adipose -tissue and decreases that of muscles - A clue for increased fat depot and insulin resistance. *Endocrinology*. 1990; 127: 3246-3248.
 11. Diab A, Dastmalchi LN, Gulati M, Michos ED. A Heart-Healthy Diet for Cardiovascular Disease Prevention: Where Are We Now? *Vasc Health Risk Manag*. 2023; 19: 237-253. doi: 10.2147/VHRM.S379874.
 12. Macknin M, Kong T, Weier A, Worley S, Tang AS, Alkhoury N et al. Plant-based, no-added-fat or American Heart Association diets: impact on cardiovascular risk in obese children with hypercholesterolemia and their parents. *J Pediatr*. 2015; 166(4): 953-9.e1-3. doi: 10.1016/j.jpeds.2014.12.058. Epub 2015 Feb 12.
 13. Luna-Castillo KP, Olivares-Ochoa XC, Hernández-Ruiz RG, Llamas-Covarrubias IM, Rodríguez-Reyes SC, Betancourt-Núñez A et al. The Effect of Dietary Interventions on Hypertriglyceridemia: From Public Health to Molecular Nutrition Evidence. *Nutrients*. 2022; 14(5): 1104. doi: 10.3390/nu14051104.
 14. Samuel PO, Edo GI, Emakpor OL, Oloni GO, Ezekiel GO, Essaghah AEA et al. Lifestyle modifications for preventing and managing cardiovascular diseases. *Sport Sci Health*. 2024; 20(1): 23-36.
 15. Khan MI, Sameen A, eds. *Animal Sourced Foods for Developing Economies: Preservation, Nutrition, and Safety*. CRC Press, 2018.
 16. Zaid M, Hasnain S. Plasma lipid abnormalities in Pakistani population: trends, associated factors, and clinical implications. *Braz J Med Biol Res*. 2018; 51(9): e7239. doi: 10.1590/1414-431X20187239.
 17. Thomas MS, Calle M, Fernandez ML. Healthy plant-based diets improve dyslipidemias, insulin resistance, and inflammation in metabolic syndrome. A narrative review. *Adv Nutr*. 2023; 14(1): 44-54. doi: 10.1016/j.advnut.2022.10.002. Epub 2022 Dec 17.
 18. Ge P, Dong C, Ren X, Weiderpass E, Zhang C, Fan H et al. The High Prevalence of Low HDL-Cholesterol Levels and Dyslipidemia in Rural Populations in Northwestern China. *PLoS One*. 2015; 10(12): e0144104. doi: 10.1371/journal.pone.0144104.
 19. West CE, Sullivan DR, Katan MB, Halferkamp IL, van der Torre HW. Boys from populations with high-carbohydrate intake have higher fasting triglyceride levels than boys from populations with high-fat intake. *Am J Epidemiol*. 1990; 131(2): 271-82. doi: 10.1093/oxfordjournals.aje.a115497.
 20. Calder PC, Deckelbaum RJ. Editorial: Omega-3 fatty acids and cardiovascular outcomes: an update. *Curr Opin Clin Nutr Metab Care*. 2019; 22(2): 97-102.
 21. Thomas MS, Calle M, Fernandez ML. Healthy plant-based diets improve dyslipidemias, insulin resistance, and inflammation in metabolic syndrome. A narrative review. *Adv Nutr*. 2023; 14(1): 44-54. doi: 10.1016/j.advnut.2022.10.002. Epub 2022 Dec 17.
 22. Espírito Santo LR, Faria TO, Silva CSO, Xavier LA, Reis VC, Mota GA et al. Socioeconomic status and education level are associated with dyslipidemia in adults not taking lipid-lowering medication: a population-based study. *Int Health*. 2022; 14(4): 346-353. doi: 10.1093/inthealth/ihz089.

