

Association of Dry Eye Disease and Diabetic Retinopathy with Glycated Hemoglobin at a Tertiary Care Unit of Karachi Pakistan

Sharjeel Sultan¹, Mahtab Alam Khanzada², Ayisha Shakeel³, Nida Ahmed⁴,
Asher Fawwad^{4*}, Abdul Basit⁵

ABSTRACT

OBJECTIVE: To investigate the association of dry eye disease and diabetic retinopathy (DR) with glycated haemoglobin at a tertiary care unit in Karachi, Pakistan.

METHODOLOGY: This cross-sectional observational study was conducted at Baqai Institute of Diabetology and Endocrinology, Baqai University Karachi, Pakistan, from July to December 2020. A total of 238 subjects having type 2, type 1, and gestational diabetes mellitus participated in this study. They went through a routine ophthalmic examination, breakup tear film time (BUT) test, Schirmer I test, staining fluorescein, and fundus photography performed to diagnose the DR. Baseline detail and biochemical parameters were recorded. Data analysis was done on statistical packages for the sciences social (SPSS) 20 version.

RESULTS: A total of 72 participants had dry eye disease (DED) conforming to a total prevalence of 30.25%; 11(4.6%) had severe dry eyes, 26(10.9%) had moderate dry eyes, 35(14.7%) had mild dry eyes whereas, 166(69.7%) participants had normal eyes. There was no significant association between gender, index mass body (BMI), smoking habits, history of family diabetes, and duration of Diabetes with DED. The frequency of diabetic retinopathy (DR) was registered as 23.5%; 29(24%) males and 27 (23.1%) females, respectively.

CONCLUSION: Overall, a 30.3% frequency of dry eye in diabetic individuals was observed. It should improve by having consistent follow-ups after three to six months, providing a distinct difference in the condition compared to the non-affected individuals.

KEYWORDS: Dry eyes, ocular surface disease, Dryness in the eyes, deficiency tear, lacrimal function unit, Schirmer's test, Diabetes.

INTRODUCTION

Worldwide, Diabetes is a rising trend, and it is projected by the Diabetes International Federation (IDF) that around 537 million folks are surviving with Diabetes, with predictions expected to upsurge above 643 million individuals by 2030¹. In Pakistan, Diabetes is now known as a chronic disease due to the adaptation of Westernized diets and various changes in lifestyle and demographic features. According to the recent second National Pakistan Diabetes Survey (NDSP) 2016-2017, Diabetes has affected 26% of the country's population, which is alarming and a

significant public health issue². Diabetes, if not under reasonable control, can cause chronic complications that include retinopathy, nephropathy, neuropathy, and heart and vascular diseases.³

Blindness's major cause is DR in developed and underdeveloped countries¹². The prevalence of diabetic retinopathy (DR) in Pakistan was 28.78%⁴. DR itself is a leading cause of blindness, and dry eye disease (DED) aggravates vision loss which usually may lead to poor quality of life⁵. DED has corneal complications such as superficial keratopathy punctuates, trophic ulceration, tear film hyperosmolarity, instability, neurosensory abnormalities, and persistent epithelial defects⁶. Hyperosmolarity and instability tear film are the procedures that cause dry eyes with Diabetes⁷. Tear film abnormality is a significant diabetic feature of the ocular disease surface due to low quality and tear functions that occur with the subnormal ocular surface⁸.

Dry eyes are commonly observed in the age group between 60-70 years⁹. It's because aging results in acinar degeneration and nuclear abnormalities that entirely change the Lacrimal unit's function. It has also been reported that male gender, prolonged Diabetes, and elevated glycated haemoglobin (HbA1c) levels were positively linked to greater severity of DR¹⁰. A

¹Department of Ophthalmology, Dow University of Health Sciences, Karachi-Pakistan

²Institute of Ophthalmology, Liaquat University of Medical & Health Sciences, Jamshoro

³Al Nafees Medical College and Hospital, Isra University, Islamabad-Pakistan

⁴Baqai Institute of Diabetology and Endocrinology, Baqai Medical University, Karachi-Pakistan

⁵Department of Medicine, Baqai Medical University, Karachi-Pakistan

*Correspondence: asherfawwad@bide.edu.pk

doi: 10.22442/jlumhs.2023.00977

Received: 18-07-2022

Revised: 03-04-2023

Accepted: 10-04-2023

Published Online: 14-04-2023



direct relationship was also found in an investigation between HbA1c levels and the prevalence of DED. The greater the HbA1c levels, the more likely to acquire dry eye conditions⁶.

Early diagnosis of dry eyes has now become necessary for clinicians to determine the associated factors and to prevent visual acuity with early treatment. With the scarcity of data from this globe, we aim to assess the association of dry eye disease and diabetic retinopathy (DR) with glycated haemoglobin at a tertiary care unit in Karachi, Pakistan.

METHODOLOGY

This cross-sectional study was conducted at Baqai Institute of Diabetology and Endocrinology (BIDE), Medical Baqai University (BMU), Karachi – Pakistan. The study period was from July to December 2020; ethical approval was obtained from BIDE's Institutional Review Board (IRB). Each study subject was given a detailed methodology description, and written and verbal consent was obtained. Both males and females with known type 2 diabetes mellitus (DM), type I DM, and gestational DM, aged between 35-70 years, were included. Subjects with other types of DM and those taking other than Diabetes medications which can affect the production of tears were omitted from the study. Subjects with previous ocular surgery history, palsy Bell's person, Sjogren's syndrome, rheumatoid arthritis, or Parkinson's disease were also excluded.

Type II DM subjects were selected using a random sampling technique. Baseline details were obtained on a predesigned questionnaire. It includes diabetes duration, types of Diabetes, gender, age, smoking habit, alcohol habit, Diabetes family history, height, weight and blood pressure. Recent haemoglobin (Hb) A1c value and lipid profile levels were recorded from BIDE's hospital management system (HMS). A routine ophthalmic examination was done, which included a tear film breakup time (TBUT) test, Schirmer test, fluorescein staining and fundus photography for DR. Dry eyes were suspected based on a history of ocular discomforts such as soreness, gritty sensation, itchiness, redness, excessive watering of the eye, and blurred vision that improve after blinking. Subjects underwent lamp slit examination of surface ocular staining dye pattern with stain fluorescein. For the confirmation of the condition, a strip Schirmer test was made. The strip was placed at the middle junction and the third lateral of the lower eyelid, and the amount of wetting was measured after 5 minutes. The condition was graded on the scale: 0-5 mm; severe dry eye, 5-10 mm; moderately dry eye, 10-15 mm; mild dry eye, and ≥ 15 mm; normal tear function¹¹.

The glycemic index was targeted as HbA1c <7% (good glycemic control) and >7% (bad glycemic control)¹². The subject was considered dyslipidemic if having a serum total cholesterol >200 mg/dl, serum LDL- cholesterol >130 mg/dl, serum HDL- cholesterol <40 mg/dl (for males) and <50 mg/dl (for females), and serum triglycerides >150 mg/dl¹³. Height was

measured by directly measuring the length from the bottom of the feet to the highest point of the head. Weight was measured in kilograms by a weighing machine. BMI was calculated by dividing weight (kg)/ height (m)². BMI <23 kg/m² was considered normal, ≥ 23 -24.9 kg/m² overweight and ≥ 25 kg/m² obese per Asia Pacific Guideline¹⁴. Blood pressure was measured in a sitting position after 10 minutes of rest using a mercury sphygmomanometer. Hypertension was considered if the subject had blood pressure $\geq 140/90$ mmHg¹⁵.

Statistical analysis

Data was analyzed on Statistical Packages for Social Sciences (SPSS) version 20. Continuous variables were presented as mean \pm standard deviation (SD), while categorical variables were presented as numbers (percentages). The chi-square, ANOVA, and Kruskal-Wallis tests were applied where applicable to determine the association between variables.

RESULTS

A total of 238 individuals were assessed. **Table I**

Table I: Baseline characteristics of studied subjects

| Variables | N (%) or Mean \pm S.D | |
|--------------------------------------|-------------------------|------------|
| N | 238 | |
| Gender | Male | 121(50.8%) |
| | Female | 117(49.2%) |
| Age (years) | 50.52 \pm 11.85 | |
| Marital status | Single | 13(5.5%) |
| | Married | 214(90.7%) |
| | Widow | 9(3.8%) |
| Body mass index (kg/m ²) | 28.94 \pm 5.28 | |
| Systolic blood pressure (mm Hg) | 124.15 \pm 18.51 | |
| Diastolic blood pressure (mmHg) | 77.43 \pm 9.41 | |
| Type of Diabetes | Type 1 | 3(1.3%) |
| | Type 2 | 234(98.3%) |
| | Gestational | 1(0.4%) |
| Smoking habit | No | 169(81.3%) |
| | Ex-smoker | 19(9.1%) |
| | Current smoker | 20(9.6%) |
| Alcohol addiction | No | 206(99%) |
| | Yes | 2(1%) |
| Family history of Diabetes | No | 51(24.5%) |
| | Yes | 157(75.5%) |
| HbA1c (%) | 9.13 \pm 1.99 | |
| Cholesterol (mg/dl) | 161.28 \pm 44.76 | |
| Triglyceride(mg/dl) | 191.24 \pm 126.53 | |
| High density lipoprotein(mg/dl) | 32.75 \pm 8.24 | |
| Low-density lipoprotein (mg/dl) | 112.15 \pm 39.87 | |
| Duration of DM | ≤ 5 years | 110(46.2%) |
| | 5 to 10 years | 57(23.9%) |
| | >10 years | 71(29.8%) |

shows the mean age of subjects (121 males, 117 females) was 50.5 ± 11.85 years. The study subjects were primarily non-smokers, 169(81.3%), and married 214(90.7%). Cholesterol total, triglycerides, HDL, and LDL, were 161.28 ± 44.76 , 191.24 ± 126.53 , 32.75 ± 8.24 , and 112.15 ± 39.87 , respectively. Poor glycemic control was observed with mean HbA1c levels of 9.13 ± 1.99 . More than half of the subjects, 157(75.5%), had a family history of Diabetes, and 110 (46.2%) participants had Diabetes for less than five years.

Table II represents the assessment of tear meniscus height using Schirmer's strip. Out of 238 subjects, 11 (4.6%) had severe dry eyes, 26(10.9%) had moderate dry eyes, 35(14.7%) had mild dry eyes whereas, 166 (69.7%) individuals had normal eyes. The overall frequency of dry eye in diabetic individuals was found to be 72(30.25%).

Table II: Assessment of tear meniscus height using Schirmer's strip method

| Schirmer's strip test (mm) | n (%) | Overall |
|----------------------------|------------|------------|
| N | 238 | |
| ≤5 | 11(4.6%) | |
| 6-10 | 26(10.9%) | 72(30.25%) |
| 11-15 | 35(14.7%) | |
| >15 | 166(69.7%) | |

In males, the association was found to be 63.6% for severe, 53.8% for moderate, and 40% for mild dry eyes, whereas, in females, the association was 36.4%, 46.2%, and 60%, respectively. All the individuals lie in the obese range of ≥ 25 kg/m² for normal to severe dry eyes. Individuals with a positive history of Diabetes and a duration of fewer than five years suffered from moderate to severe dry eyes. There was no significant association between gender, body mass index (BMI), smoking habits, family history of Diabetes, and duration of Diabetes with tear breakup time (TBUT), p-value <0.05. Upon evaluating the relationship between biochemical parameters with TBUT, poor glycemic control was observed in severe (8.87 ± 1.04), mild (9.45 ± 2.31), and moderate dry eyes (9.84 ± 1.33). No significant association was found between lipid profile, HbA1c, and TUBT, while a significant association was found in cholesterol levels among severe (170 ± 53.39), mild (186.86 ± 59.23), moderate (173.94 ± 46.93), and normal eyes (152.83 ± 38.58) p-value = 0.03 (**Table III**).

In participants who have Diabetes, the prevalence of diabetic retinopathy was found to be 23.5%. Among 56(23.5%) individuals having retinopathy, 29(24%) were males, whereas 27(23.1%) were females (**Table IV**). Furthermore, **Table IV** and **Table V** show the association of diabetic retinopathy with dry eye

Table III: Association of tear film breakup time (TBUT) with baseline and biochemical parameters

| Variable | TBUT (mm) | | | | P-value | |
|---|--------------------|--------------------|--------------------|--------------------|------------|-------|
| | ≤5 | 6-10 | 11-15 | >15 | | |
| N | 11 | 26 | 35 | 166 | | |
| Age (years) | 47.91 ± 12.58 | 48.54 ± 11.75 | 51.43 ± 11.24 | 50.81 ± 11.99 | 0.868 | |
| Gender | Male | 7(63.6%) | 14(53.8%) | 14(40%) | 86(51.8%) | 0.471 |
| | Female | 4(36.4%) | 12(46.2%) | 21(60%) | 80(48.2%) | |
| Systolic blood pressure (mm Hg) | 137.14 ± 33.02 | 123.78 ± 19.16 | 124.34 ± 15.95 | 123.54 ± 18.03 | 0.77 | |
| Diastolic blood pressure (mmHg) | 77.14 ± 11.13 | 78.87 ± 9.16 | 77.16 ± 8.8 | 77.28 ± 9.57 | 0.895 | |
| Body mass index (kg/m²) | 30.05 ± 3.95 | 29.42 ± 4.79 | 28.2 ± 4.19 | 28.97 ± 5.65 | 0.772 | |
| Family history of Diabetes | No | 2(28.6%) | 5(21.7%) | 10(31.2%) | 34(23.3%) | 0.786 |
| | Yes | 5(71.4%) | 18(78.3%) | 22(68.8%) | 112(76.7%) | |
| Duration of Diabetes | ≤5 years | 6(54.5%) | 12(46.2%) | 9(25.7%) | 83(50%) | 0.114 |
| | 5-10 years | 3(27.3%) | 9(34.6%) | 11(31.4%) | 34(20.5%) | |
| | >10 years | 2(18.2%) | 5(19.2%) | 15(42.9%) | 49(29.5%) | |
| Type of Diabetes | Type 2 | 11(100%) | 26(100%) | 35(100%) | 162(97.6%) | 0.623 |
| | Other | 0(0%) | 0(0%) | 0(0%) | 4(2.4%) | |
| HbA1c (%) | 8.87 ± 1.04 | 9.45 ± 2.31 | 9.84 ± 1.33 | 8.93 ± 2.06 | 0.275 | |
| Cholesterol (mg/dl) | 170 ± 53.39 | 186.86 ± 59.23 | 173.94 ± 46.93 | 152.83 ± 38.58 | 0.032 | |
| Triglyceride(mg/dl) | 250.5 ± 238.45 | 176.07 ± 64.43 | 211.94 ± 128.6 | 186.01 ± 128.9 | 0.717 | |
| High-density lipoprotein (mg/dl) | 31.75 ± 6.13 | 35.14 ± 7.44 | 36.18 ± 7.54 | 31.53 ± 8.45 | 0.122 | |
| Low-density lipoprotein (mg/dl) | 98.25 ± 40.94 | 123.67 ± 53.73 | 121.05 ± 35.46 | 108.64 ± 38.08 | 0.432 | |

Data presented as mean \pm SD or n (%); P-value <0.05 considered statistically significant.

disease and gender. In contrast, no association significantly was found between diabetic retinopathy, dry eye disease, and gender was found.

Table IV: Association of Retinopathy with TBUT

| Retinopathy | TBUT (mm) | | | | P-value |
|-------------|--------------|---------------|---------------|----------------|---------|
| | ≤5 | 6-10 | 11-15 | >15 | |
| No# | 11 | 26 | 35 | 166 | |
| No | 9 (81.8%) | 21 (80.8%) | 26 (74.3%) | 126 (75.9%) | 0.905 |
| Yes | 2 (18.2%) | 5(19.2%) | 9(25.7%) | 40 (24.1%) | |

Data presented as n (%); P-value<0.05 considered as statistically significant; NO# = numbers

Table V: Association of retinopathy with gender

| Retinopathy | Male | Female | P-value | Overall |
|-------------|------------|------------|----------|------------|
| N | 121 | 117 | - | 238 |
| No | 92(76%) | 90(76.9%) | 0.871 | 182(76.5%) |
| Yes | 29(24%) | 27(23.1%) | | 56(23.5%) |

Data presented as n (%); P-value<0.05 considered as statistically significant

DISCUSSION

Overall, a 30.3% frequency of dry eye in diabetic individuals was observed, among which 4.6% had severe dry eyes, 10.9% had moderate dry eyes, and 14.7% had mild dry eyes. The prevalence of diabetic retinopathy was 23.5%, and participants were observed with poor glycemic control and bad triglyceride levels. The DED in diabetic patients varies in different populations. Fuerst N et al.¹⁶ found a 47.8% dry eye prevalence higher than our study.

Dry eyes were more frequent in females than males, in line with a recent study that showed females were more prone to dry eye disease than males¹⁷. Females were more inclined to males, which may be due to the onset of menopause that causes decreased estrogen levels in females and leads to reduced tear film⁹. Misra SL et al.¹⁸ reported that due to the lesser production of androgens, a type of protective hormone, there is an upsurge in the incidence of dry eye disease in women. However, other studies have negated the association of gender in diabetic individuals with dry eye disease¹⁹. This research revealed that reduced tear formation in some diabetic individuals is linked to autonomic nervous system dysfunction²⁰. We found the component of total cholesterol were similar in levels among diabetic participants with DED and non-DED. Our data show no association between dry eye disease and lipid profile. In our study, the prevalence of diabetic retinopathy was 23%, and gender had no statistically significant influence on the condition. We observed most participants with less than five years of Diabetes,

in contrast with ul Islam Q 2017²¹ studies which found participants with longer diabetes duration.

In this investigation, we observed participants with poor glycemic control; however, the association of HbA1c with dry eye was statistically insignificant, in contrast with the findings of other studies²²⁻²⁶, which found a strong positive connection between HbA1c and dry eye. Manjula TR 2019²⁷ stated that DM management with dry eyes is statistically significant. A study from the Asian population also found a strong association between dry eyes with poor glycemic control²⁸.

CONCLUSION

Overall, a 30.3% frequency of dry eye in diabetic individuals was observed, among which 4.6% had severe dry eyes, 10.9% had moderate dry eyes, and 14.7% had mild dry eyes. It may improve by having consistent follow-ups after three to six months, providing a distinct difference in the condition compared to the non-affected individuals.

Limitations

This study has some limitations due to its relatively small sample size and the fact that it was conducted at only one clinical site. Moreover, menopausal history, which significantly affects the development of DED, was lacking in our study.

Ethical Permission: Baqai Medical University, Karachi, IRB letter No. BIDE/IRB/SSULTAN/08/20/0234.

Conflict of Interest: No conflicts of interest.

Financial Disclosure / Grant Approval: No funding agency was involved in this research.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publically.

AUTHOR CONTRIBUTIONS

Sultan S: Concept, design, edited and approved the manuscript

Khanzada MA: Interpretation of data, edited and approved the manuscript

Shakeel A: Interpretation of data, edited and approved the manuscript

Ahmed N: Literature search, interpretation of data, and wrote the manuscript

Fawwad A: Concept, design, Edit, and support the final manuscript

Basit A: Concept, design, Edit, and approve the final manuscript

REFERENCES

1. Federation ID. Diabetes facts & figures. International Diabetes Federation. Available at: Available at: <https://idf.org> › About Diabetes ›

- What is Diabetes. [Last accessed on 12-Sept-2021].
2. Basit A, Fawwad A, Qureshi H, Shera AS, NDSP Members. Prevalence of Diabetes, pre-diabetes and associated risk factors: Second National Diabetes Survey of Pakistan (NDSP), 2016–2017. *BMJ Open*. 2018; 8: e020961. doi: 10.1136/bmjopen-2017-020961.
 3. Chou CY, Hsu DY, Chou CH. Predicting the Onset of Diabetes with Machine Learning Methods. *J Pers Med*. 2023; 13(3): 406. doi: 10.3390/jpm13030406.
 4. Mumtaz SN, Fahim MF, Arslan M, Shaikh SA, Kazi U, Memon MS. Prevalence of diabetic retinopathy in Pakistan; A systematic review. *Pak J Med Sci*. 2018; 34(2): 493-500. doi: 10.12669/pjms.342.13819.
 5. Zou X, Lu L, Xu Y, Zhu J, He J, Zhang B et al. Prevalence and clinical characteristics of dry eye disease in community-based type 2 diabetic patients: the Beixinjing eye study. *BMC Ophthalmol*. 2018; 18(1): 117. doi: 10.1186/s12886-018-0781-7.
 6. Midhuna K, Divya N, Veeramani PA, Bhaskaran B. Assessment of the incidence and prevalence of dry eyes in diabetic individuals and its effect on visual acuity. *Int J Curr Pharm Res*. 2020; 12(6): 97-99. doi: 10.22159/ijcpr.2020v126.40301.
 7. Zhang X, Zhao L, Deng S, Sun X, Wang N. Dry eye syndrome in patients with diabetic mellitus: prevalence, etiology and clinical characteristics. *J Ophthalmol*. 2016; 2016. doi: 10.1155/2016/8201053. Epub 2016 Apr 26.
 8. Kesarwani D, Rizvi SWA, Khan AA, Amitava AK, Vasenwala SM, Siddiqui Z. Tear film and ocular surface dysfunction in Diabetes mellitus in an Indian population. *Indian J Ophthalmol*. 2017; 65(4): 301-4. doi: 10.4103/ijo.IJO_939_15.
 9. Kamel SS, Mohammed TH, El Zankalony YA, Saad AH. Prevalence of dry eyes in diabetics. *J Egyptian Ophthalmol*. 2017; 110: 77-82.
 10. Long M, Wang C, Liu D. Glycated hemoglobin A1C and vitamin D and their association with diabetic retinopathy severity. *Nutr Diabetes*. 2017; 7(6): e281. doi: 10.1038/nutd.2017.30.
 11. Mohmond AJ. Treatment of Dry Eye: comparing the use of Tears Naturale Eye Drop, Omega 3, Physical Treatment and Exercises (Doctoral dissertation, Alneelain University). 2014.
 12. Fawwad A, Waris N, Basit KA, Tahir B, Gregg EW, Basit A. NDSP-10: The Cardiometabolic Risk Profile with Various Degrees of Dysglycemia in Younger and Older Adults: Findings from the Second National Diabetes Survey of Pakistan 2016–2017. *Metab Syndr Relat Disord*. 2022; 20(6): 351-59. doi: 10.1089/met.2021.0154. Epub 2022 May 26.
 13. Basit A, Sabir S, Riaz M, Fawwad A, NDSP Members. NDSP 05: Prevalence and pattern of dyslipidemia in urban and rural areas of Pakistan; A sub-analysis from second National Diabetes Survey of Pakistan (NDSP) 2016–2017. *J Diabetes Metab Disord*. 2020; 19(2): 1215-25. doi: 10.1007/s40200-020-00631-z.
 14. Pan WH, Yeh WT. How to define obesity? Evidence-based multiple action points for public awareness, screening, and treatment: an extension of Asian-Pacific recommendations. *Asia Pac J Clin Nutr*. 2008; 17(3): 370-4.
 15. Basit A, Tanveer S, Fawwad A, Naeem N, NDSP Members. Prevalence and contributing risk factors for hypertension in urban and rural areas of Pakistan; a study from second National Diabetes Survey of Pakistan (NDSP) 2016–2017. *Clin Exp Hypertens*. 2020; 42(3): 218-24. doi: 10.1080/10641963.2019.1619753. Epub 2019 May 31.
 16. Fuerst N, Langelier N, Massaro-Giordano M, Pistilli M, Stasi K, Burns C et al. Tear osmolarity and dry eye symptoms in diabetics. *Clin Ophthalmol*. 2014; 8: 507-15. doi: 10.2147/OPHTH.S51514.
 17. Beckman KA. Characterization of dry eye disease in diabetic patients versus non-diabetic patients. *Cornea*. 2014; 33(8): 851-4. doi: 10.1097/ICO.000000000000163.
 18. Misra SL, Craig JP, Patel DV, McGhee CN, Pradhan M, Ellyett K et al. In vivo confocal microscopy of corneal nerves: an ocular biomarker for peripheral and cardiac autonomic neuropathy in type 1 diabetes mellitus. *Invest Ophthalmol Vis Sci*. 2015; 56(9): 5060-5. doi: 10.1167/iovs.15-16711.
 19. Achtsidis V, Eleftheriadou I, Kozanidou E, Voumvourakis K, Stamboulis E, Theodosiadis PG et al. Dry eye syndrome in subjects with Diabetes and association with neuropathy. *Diabetes Care*. 2014; 37(10): e210-1. doi: 10.2337/dc14-0860.
 20. Ramos-Remus C, Suarez-Almazor M, Russell AS. Low tear production in patients with Diabetes mellitus is not due to Sjögren's syndrome. *Clin Exp Rheumatol*. 1994; 12(4): 375-380.
 21. ul Islam Q, Mehboob MA, Amin ZA. Comparison of corneal morphological characteristics between diabetic and nondiabetic population. *Pak J Med Sci*. 2017; 33(6): 1307-1311. doi: 10.12669/pjms.336.13628.
 22. Sagdik HM, Ugurbas SH, Can M, Tetikoglu M, Ugurbas E, Ugurbas SC et al. Tear film osmolarity in patients with diabetes mellitus. *Ophthalmic Res*. 2013; 50(1): 1-5. doi: 10.1159/000345770. Epub 2013 Feb 22.

23. He F, Zhao Z, Liu Y, Lu L, Fu Y. Assessment of Ocular Surface Damage during the Course of Type 2 Diabetes Mellitus. *J Ophthalmol*. 2018; 2018: 1206808. doi:10.1155/2018/1206808.
24. Nadeem H, Malik TG, Mazhar A, Ali A. Association of Dry Eye Disease with Diabetic Retinopathy. *J Coll Physicians Surg Pak*. 2020; 30 (5): 493-7. doi:10.29271/jcsp.2020.05.493.
25. Khetwani D, Karambelkar VH, Gadre GA. Ocular surface disorders in type 2 diabetes mellitus. *Int J Contemp Med Res*. 2019; 6(3): C9-C12. doi: 10.21276/ijcmr.2019.6.3.41.
26. Najafi L, Malek M, Valojerdi AE, Aghili R, Khamseh ME, Fallah AE et al. Dry eye and its correlation to Diabetes microvascular complications in people with type 2 diabetes mellitus. *J Diabetes Complications*. 2013; 27(5): 459-62. doi: 10.1016/j.jdiacomp.2013.04.006. Epub 2013 May 30.
27. Manjula TR, Gahana K, Harsha R. A Clinical Study on Meibomian Gland Dysfunction and Dry Eye in Patients with Type2 Diabetes Mellitus. *J Med Sci Health*. 2019; 5(3): 7-12.
28. Waris SA, Balaji RSN, Huda R. To study prevalence of dry eyes in diabetic patients. *Indian J Clin Exp Ophthalmol*. 2019; 5: 40-43. doi: 10.18231/2395-1451.2019.0010.

